

PROVENGE[®]
(sipuleucel-T)

ACTIVATE YOUR IMMUNE SYSTEM TO FIGHT ADVANCED
PROSTATE CANCER WITH **PROVENGE**

Think **more** about **special occasions**.
Think **less** about **cancer**.
Start **PROVENGE** today!



PROVENGE extends life for certain men with metastatic CRPC. According to an exploratory analysis, men who received PROVENGE with a PSA less than or equal to 22.1 ng/mL when they started treatment saw the greatest improvement.

✓ Increased median overall survival of over 1 year*


The most common side effects of PROVENGE are generally mild to moderate and well tolerated.

The PROVENGE treatment process can be completed in 6 appointments throughout the course of about a month.

*Median overall survival improvement was found in all quartiles of the analysis (PSA ≤22.1 ng/mL: 13 months, PSA >22.1-50.1 ng/mL: 7.1 months, PSA >50.1-134.1 ng/mL: 5.4 months, PSA >134.1 ng/mL: 2.8 months) Individual results may vary.

See back page for Indication and Important Safety Information.

Visit **www.PROVENGE.com** for more details



PROVENGE®

(sipuleucel-T)

INDICATION

PROVENGE® (sipuleucel-T) is a cellular immunotherapy used to treat certain men with advanced prostate cancer. PROVENGE is made from your own immune cells.

IMPORTANT SAFETY INFORMATION

Before getting PROVENGE, tell your doctor about all your medical problems, including problems with your heart or lungs, or if you have had a stroke.

Tell your doctor about all the medicines you take, including prescription and nonprescription drugs, vitamins, and dietary supplements.

PROVENGE is made from your own immune cells, collected approximately 3 days before each scheduled infusion of PROVENGE. The collection is called “leukapheresis” (pronounced loo-kuh-fuh-REE-sis). You will get PROVENGE in 3 intravenous infusions (put into your veins), about 2 weeks apart. Each infusion takes about 60 minutes. You will then be monitored for at least 30 minutes. Your doctor will give you a schedule for your cell collection and infusion appointments. It is very important that you arrive on time for your appointments.

The most common side effects of PROVENGE include chills, fatigue, fever, back pain, nausea, joint ache, and headache. These are not all the possible side effects of PROVENGE treatment.

PROVENGE infusion can cause serious reactions. Tell your doctor right away if you:

- Have breathing problems, chest pains, racing heart or irregular heartbeats, high or low blood pressure, dizziness, fainting, nausea, or vomiting after getting PROVENGE. Any of these may be signs of heart or lung problems.
- Develop numbness or weakness on one side of the body, decreased vision in one eye, or difficulty speaking. Any of these may be signs of a stroke.
- Develop symptoms of thrombosis which may include: pain and/or swelling of an arm or leg with warmth over the affected area, discoloration of an arm or leg, unexplained shortness of breath, chest pain that worsens on deep breathing.
- Get a fever over 100°F, or redness or pain at the infusion or collections sites. Any of these may be signs of infection.

Tell your doctor about any side effect that concerns you or does not go away. For more information, talk with your doctor.

You are encouraged to report negative side effects of prescription drugs to the FDA.
Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see enclosed full Prescribing Information.

